

AN EQUIVALENT OF KRONECKER'S THEOREM FOR POWERS OF AN ALGEBRAIC NUMBER AND STRUCTURE OF LINEAR RECURRENCES OF FIXED LENGTH

NEVIO DUBBINI AND MAURIZIO MONGE

ABSTRACT. After defining a notion of ϵ -density, we provide for any integer $m > 1$ and real algebraic number α an estimate of the smallest ϵ such that the set of vectors of the form $(t, t\alpha, \dots, t\alpha^{m-1})$ for $t \in \mathbb{R}$ is ϵ -dense modulo 1 in terms of the multiplicative Mahler measure $M(A(x))$ of the minimal integral polynomial $A(x)$ of α , which is independent of m . In particular, we show that if α has degree d it is possible to take $\epsilon = 2^{\lfloor d/2 \rfloor} / M(A(x))$.

On the other side we show using asymptotic estimates for Toeplitz determinants that we cannot have ϵ -density for sufficiently large m whenever ϵ is strictly smaller than $1/M(A(x))$. In the process of proving this we obtain a result of independent interest about the structure of the $\mathbb{Z}$ -module of integral linear recurrences of fixed length determined by a non-monic polynomial.

1. INTRODUCTION

The classical Kronecker's Theorem in diophantine approximation, in one of its different versions, says that if $\theta_1, \dots, \theta_m$ are real numbers linearly independent over $\mathbb{Q}$, then $(t\theta_1, \dots, t\theta_m)$ is dense modulo 1, i.e. if we call

$$\pi_m : \mathbb{R}^m \rightarrow \mathbb{R}^m / \mathbb{Z}^m = \mathbb{T}^m$$

the canonical projection, then the set $\{\pi_m(t\theta_1, \dots, t\theta_m), t \in \mathbb{R}\}$ is dense in $\mathbb{T}^m$.

A question left to investigate by this result is which amount it fails for particular choices of (θ_i) which are not linearly independent over $\mathbb{Q}$, but have a special form. In this paper we will consider the case where the (θ_i) are powers of an algebraic number α , and more in general when the θ_i satisfy a linear recurrence relation with characteristic polynomial $A(x)$ of degree d , and $\theta_1, \theta_2, \dots, \theta_d$ are linearly independent over $\mathbb{Q}$ (this condition is automatically satisfied when $\theta_i = \alpha^{i-1}$, taking the minimal integral polynomial of α as $A(x)$).

Before stating our main result, we shall give a definition of ϵ -density and recall the notion of Mahler measure.

Date: February 8, 2010.

2000 *Mathematics Subject Classification.* Primary: 11K06, 11K60, Secondary: 11G50, 11H31.

1.1. **ϵ -density.** For $\epsilon > 0$ and positive integer m , let

$$C_\epsilon = \pi_m(I_\epsilon) \subseteq \mathbb{T}^m, \quad \text{where } I_\epsilon = \left[-\frac{\epsilon}{2}, \frac{\epsilon}{2}\right]^m,$$

be the cube with edge length ϵ in the torus $\mathbb{T}^m$, which we assume centered around the origin for convenience.

Definition. We will say that a set $S \subseteq \mathbb{T}^m$ is ϵ -dense if $S + C_{\bar{\epsilon}} = \mathbb{T}^m$ for each $\bar{\epsilon} > \epsilon$, or equivalently if $S + C_\epsilon$ is dense. A set $T \in \mathbb{R}^m$ will be ϵ -dense if $T + I_{\bar{\epsilon}} = \mathbb{R}^m$ whenever $\bar{\epsilon} > \epsilon$, or equivalently if $T + I_\epsilon$ is dense.

Alternatively, it is possible to consider on $\mathbb{T}^m$ the distance

$$d_\infty(x, y) = \inf \left\{ |\tilde{x} - \tilde{y}|_\infty, \tilde{x}, \tilde{y} \in \mathbb{R}^m, \pi_m(\tilde{x}) = x, \pi_m(\tilde{y}) = y \right\},$$

and since the $\rho/2$ -neighborhood of S with respect to $d_\infty(\cdot, \cdot)$ is precisely $S + C_\rho$ for each $\rho > 0$, we have that S will be ϵ -dense if and only if its $\bar{\epsilon}/2$ -neighborhood is all of $\mathbb{T}^m$, for each $\bar{\epsilon} > \epsilon$ (or, equivalently, if its $\epsilon/2$ -neighborhood is dense).

1.2. **Mahler measure.** Suppose given a polynomial of degree d with complex coefficients

$$A(x) = \sum_{i=0}^d a_i x^i = a_d \cdot \prod_{i=1}^d (x - \alpha_i),$$

and such that $a_0 \neq 0$. The *Mahler measure* of $A(x)$ is defined as

$$M(A) = M(A(x)) = |a_d| \cdot \prod_{i=1}^d \max \{1, |\alpha_i|\}.$$

The Mahler measure is a notion of complexity that for the minimal polynomial of a rational number p/q with $(p, q) = 1$ reduces to $\max(|p|, |q|)$. Furthermore, each coefficient a_i can be written as the sum of $\binom{d}{i}$ monomials with coefficient 1 in $a_d, \alpha_1, \dots, \alpha_d$, each appearing at most once in each monomial, and consequently we have that $a_i \leq \binom{d}{i} \cdot M(A)$, for each $i = 0, \dots, d$.

Note also that

$$\begin{aligned} M(A(x/2)) &= \frac{|a_d|}{2^d} \cdot \prod_{j=1}^d \max \{1, |2\alpha_j|\} \\ &= |a_d| \cdot \prod_{j=1}^d \max \left\{ \frac{1}{2}, |\alpha_j| \right\}, \end{aligned}$$

while

$$\begin{aligned} 2^{-d} M(A(2x)) &= |a_d| \cdot \prod_{j=1}^d \max \left\{ 1, \left| \frac{\alpha_j}{2} \right| \right\} \\ &= |a_0| \cdot \prod_{j=1}^d \max \left\{ |\alpha_j^{-1}|, \frac{1}{2} \right\} = M(\tilde{A}(x/2)), \end{aligned}$$

where we called $\tilde{A}(x)$ the conjugate polynomial $\sum_{i=0}^d a_i x^{d-i}$.

We have that $M(A(x/2)) \geq 2^{-[d/2]} M(A(x))$ if $A(x)$ has at least $d/2$ roots ≥ 1 in absolute value, while $2^{-d} M(A(2x)) \geq 2^{-[d/2]} M(A(x))$ if $A(x)$ has at least $d/2$ roots ≤ 1 in absolute value. Since at least one of these conditions must hold, we have

$$\min \left\{ \frac{1}{M(A(x/2))}, \frac{2^d}{M(A(2x))} \right\} \leq \frac{2^{[d/2]}}{M(A(x))}.$$

1.3. Main results. We can now state our main results:

Theorem 1. *Let $m \geq d > 0$ be positive integers, and let $\theta_1, \dots, \theta_m$ be real numbers such that $\theta_1, \dots, \theta_d$ are linearly independent over $\mathbb{Q}$, and $\theta_{d+1}, \dots$ are inductively defined by the linear recurrence determined by a primitive integral polynomial $A(x)$ of degree d and with nonzero constant coefficient. Then*

$$S_\theta = S_{\theta, m} = \{ \pi_m(t\theta_1, t\theta_2, \dots, t\theta_m), t \in \mathbb{R} \} \subseteq \mathbb{T}^m$$

is ϵ -dense for each ϵ at least

$$\min \left\{ \frac{1}{M(A(x/2))}, \frac{2^d}{M(A(2x))} \right\} \leq \frac{2^{[d/2]}}{M(A(x))}.$$

In other terms, this theorem says that if ϵ is as stated, then for arbitrary real numbers $x_1, \dots, x_m$ it is possible to find a real number t and integers $p_1, \dots, p_m$ such that

$$x_i \leq t\theta_i - p_i \leq x_i + \epsilon, \quad \text{for } 1 \leq i \leq m.$$

It is clearly possible to take as (θ_i) the powers (α^{i-1}) of an algebraic number α , and as $A(x)$ as the minimal integral polynomial of α . For each primitive integral polynomial $A(x)$ it is also possible to take as (θ_i) any ‘sufficiently generic’ recurrence sequence determined by $A(x)$, such that $\theta_1, \dots, \theta_d$ are linearly independent over $\mathbb{Q}$.

Some computational evidence actually makes us advance the following

Conjecture 1. *In Theorem 1 we have ϵ -density for each ϵ at least $1/M(A(x))$.*

This conjectured optimal density coefficient is not very far from what obtained in Theorem 1, but it would be optimal because of the following negative result:

Theorem 2. *Let $\theta_1, \theta_2, \dots$ and $A(x)$ be as in Theorem 1. Then for each $\epsilon < 1/M(A(x))$ the set $S_{\theta, m}$ is not ϵ -dense for m big enough.*

Note that if could be ruled out the existence of $S_{\theta, m}$ which are not ϵ -dense for some $\epsilon < 1$ when $A(x)$ is not a product of cyclotomic polynomials, then a positive solution of Lehmer problem [5, Chap. 1] would follow, because the existence of a sequence of polynomials with Mahler measure approaching to 1 from above would be automatically ruled out.

During the proof of this Theorem a result about the structure of the module of linear recurrences of fixed length is obtained, it is stated as Theorem 3 after some notation has been introduced.

1.4. Acknowledgements. We wish to thank Roberto Dvornicich for offering kind support and advice during our work, and for providing us the motivation to improve the second part of the paper. We would also like to thank Albrecht Böttcher and Samuele Mongodi for their help while looking for references.

2. PRELIMINARIES

Suppose given a polynomial with complex coefficients

$$A(x) = a_d x^d + \cdots + a_1 x + a_0,$$

and suppose that a_d and a_0 are both nonzero. For each $\ell \geq 1$ let's consider the rectangular $\ell \times (\ell + d)$ matrix

$$\llbracket A \rrbracket_\ell \stackrel{\text{def}}{=} \begin{pmatrix} a_0 & a_1 & \cdots & a_d & & & \\ & a_0 & a_1 & \cdots & a_d & & \\ & & a_0 & a_1 & \cdots & a_d & \\ & & & \ddots & & \ddots & \ddots \\ & & & & a_0 & a_1 & \cdots & a_d \end{pmatrix}.$$

$\llbracket A \rrbracket_\ell$ is also the multiplication matrix for sections of power series by the polynomial $A(x)$, in the sense that if

$$F(x) = \sum_{i=0}^{\infty} f_i x^i, \quad G(x) = \sum_{i=0}^{\infty} g_i x^i$$

are complex power series such that $G(x) = A(x) \cdot F(x)$, then

$$\begin{pmatrix} g_{d+\ell-1} \\ \vdots \\ g_{d+1} \\ g_d \end{pmatrix} = \llbracket A \rrbracket_\ell \cdot \begin{pmatrix} f_{d+\ell-1} \\ \vdots \\ f_{d+1} \\ f_d \\ \vdots \\ f_0 \end{pmatrix},$$

and in fact the range of coefficients $0, 1, \dots, d + \ell - 1$ of $F(x)$ uniquely determines the coefficients $d, d + 1, \dots, d + \ell - 1$ of $G(x)$.

Moreover, it is easy to see that if $A(x)$ factors as

$$A(x) = B(x) \cdot C(x), \quad B(x) = \sum_{i=0}^s b_i x^i, \quad C(x) = \sum_{i=0}^t c_i x^i,$$

then we have

$$\llbracket A \rrbracket_\ell = \llbracket B \rrbracket_\ell \cdot \llbracket C \rrbracket_{\ell+s} = \llbracket C \rrbracket_\ell \cdot \llbracket B \rrbracket_{\ell+t}$$

as it is easy to verify, for instance observing that multiplying a power series by $A(x)$ and discarding the lowest d coefficients gives the same result as multiplying first by $C(x)$ and discarding the lowest t coefficients, and then by $B(x)$ and discarding the lowest s coefficients (and vice-versa).

For each polynomial $A(x)$ and positive integer m we will also need the lower triangular $m \times m$ matrix defined as

$$\{A\}_m \stackrel{\text{def}}{=} \begin{pmatrix} a_d & & & & \\ a_{d-1} & a_d & & & \\ \vdots & \vdots & \ddots & & \\ a_0 & a_1 & \dots & a_d & \\ & a_0 & a_1 & \dots & a_d \\ & & \ddots & \ddots & \\ & & & a_0 & a_1 & a_d \end{pmatrix}.$$

Note that $\llbracket A \rrbracket_\ell$ is embedded in $\{A\}_{\ell+d}$ as the last ℓ rows. Similarly to the previous case, we have that if $A(x) = B(x) \cdot C(x)$, then $\{A\}_m = \{B\}_m \cdot \{C\}_m$. In particular, if $C(x)$ splits in linear factors as

$$C(x) = \prod_{i=1}^t (x - \gamma_i),$$

we have that

$$\{C\}_m = \prod_{i=1}^t \{x - \gamma_i\}_m.$$

3. PROOF OF THEOREM 1

We can now prove Theorem 1.

Proof of Theorem 1. From Kronecker's Approximation Theorem [3, pg. 53-54] the set $S_{\theta,m}$ is dense in the closed subgroup of the torus of the elements satisfying the same integral relations as the $\theta_1, \dots, \theta_m$.

Let $A(x) = \sum_{i=0}^d a_d x^d = a_d \prod_{i=1}^d (x - \alpha_i)$ be the primitive integral polynomial defining the recurrence sequence $\theta_1, \theta_2, \dots$. If $k_0, \dots, k_{m-1}$ is an integral linear relation satisfied by the θ_i (in the sense that $\sum_{i=1}^m k_{i-1} \theta_i = 0$), and $K(x) = \sum_{i=0}^{m-1} k_i x^i$ is the corresponding polynomial, we can consider the remainder $R(x) = \sum_{i \geq 0} r_i x^i$ of the division of $K(x)$ by $A(x)$. It is a polynomial of degree $\leq d-1$ and its coefficients $r_0, r_1, \dots$ still define a linear relation of the θ_i . But $\theta_1, \dots, \theta_d$ are linearly independent over $\mathbb{Q}$, so $R(x)$ must be zero.

Consequently $K(x)$ must be a multiple of $A(x)$ in $\mathbb{Q}[x]$, and thus in $\mathbb{Z}[x]$ by Gauss lemma, being $A(x)$ primitive. This shows that for $\ell = m - d$ the rows of the matrix $\llbracket A \rrbracket_\ell$ generate the $\mathbb{Z}$ -module of the integral relations of $\theta_1, \dots, \theta_m$.

The closure of $S_{\theta,m}$ is thus the projection in $\mathbb{T}^m$ of the hyperplane

$$P = P(A) = \left\{ v \in \mathbb{R}^m : \llbracket A \rrbracket_\ell \cdot v = 0 \right\},$$

the kernel of the linear application $\llbracket A \rrbracket_\ell : \mathbb{R}^m \rightarrow \mathbb{R}^\ell$.

It's inverse image under the projection π_m is $P + \mathbb{Z}^m$, and since we just showed that the sublattice of $\mathbb{Z}^m$ generated by the row of the matrix $\llbracket A \rrbracket_\ell$

is the saturated sublattice of the integral linear relations of $\theta_1, \dots, \theta_m$, we have that $P + \mathbb{Z}^m$ is also equal to

$$Q = Q(A) = \left\{ v \in \mathbb{R}^m : \llbracket A \rrbracket_\ell \cdot v \in \mathbb{Z}^\ell \right\}.$$

In fact, the $\mathbb{Z}$ -linear map $\llbracket A \rrbracket_\ell : \mathbb{Z}^m \rightarrow \mathbb{Z}^\ell$ must be surjective (this can also be seen directly considering the $\ell \times \ell$ minors modulo p for each prime p [4, Lemma 2, Chap. 1], which have gcd 1 because $A(x)$ is primitive). Now, $P + \mathbb{Z}^m \subseteq Q$ clearly, and for each vector $w \in Q$ there exist a vector $z \in \mathbb{Z}^m$ such that $\llbracket A \rrbracket_\ell \cdot w = \llbracket A \rrbracket_\ell \cdot z$, and consequently the difference $v = w - z$ is in P , and we have that $w = v + z \in P + \mathbb{Z}^m$.

We are now reduced to prove that $Q(A)$ is ϵ -dense in $\mathbb{R}^m$, for each ϵ satisfying the statement of the theorem. Applying the involution

$$(x_1, x_2, \dots, x_{m-1}, x_m) \mapsto (x_m, x_{m-1}, \dots, x_2, x_1)$$

to $Q(A)$ we obtain a set which clearly has the same ϵ -density properties as $Q(A)$, and which is $Q(\tilde{A})$, where $\tilde{A}(x)$ is the conjugated polynomial $\tilde{A}(x) = a_0 x^d + \dots + a_{d-1} x + a_d$. Consequently, we can just prove that $Q(A)$ is ϵ -dense for

$$\epsilon \geq 1/M(A(x/2)) = \left(|a_d| \cdot \prod_{i=1}^d \max \left\{ \frac{1}{2}, |\alpha_i| \right\} \right)^{-1},$$

and the ϵ -density for $\epsilon \geq 2^d/M(A(2x)) = 1/M(\tilde{A}(x/2))$ will follow from the same estimate applied to $Q(\tilde{A})$.

To obtain this estimate, note first that $Q = Q(A)$ will be ϵ -dense if and only if the map $\pi_\ell \circ \llbracket A \rrbracket_\ell$ to $\mathbb{T}^\ell$ is surjective when restricted to $I_\epsilon = I_{\epsilon, m} = [-\epsilon/2, \epsilon/2]^m$. In fact, Q is ϵ -dense if and only if an arbitrary vector $v \in \mathbb{R}^m$ is contained in $Q + I_\epsilon$, and

$$\begin{aligned} &\Leftrightarrow (v + I_\epsilon) \cap Q \neq \emptyset \quad \forall v \in \mathbb{R}^m \\ &\Leftrightarrow (w + \llbracket A \rrbracket_\ell \cdot I_\epsilon) \cap \mathbb{Z}^\ell \neq \emptyset \quad \forall w \in \mathbb{R}^\ell \end{aligned}$$

applying the matrix $\llbracket A \rrbracket_\ell$ to the expression, and where we denoted $\llbracket A \rrbracket_\ell \cdot I_\epsilon$ the image of I_ϵ under the map $\llbracket A \rrbracket_\ell$. This passage must be justified because the matrix $\llbracket A \rrbracket_\ell$ clearly does not have rank m , but since Q contains *all* the vectors that are mapped to $\mathbb{Z}^\ell$ the first intersection will be non-empty whenever the second one is (the other direction being trivial).

We will now factor $A(x)$ as a product of polynomials $B(x), C(x) \in \mathbb{R}[x]$, with a consequent factorization of $\llbracket A \rrbracket_\ell$, and prove two different estimates for each of the two factors $B(x)$ and $C(x)$. We will select *a posteriori* the factorization which provides the best compound estimate.

More precisely, let $B(x) = \sum_{i=0}^s b_i x_i$ and $C(x) = \sum_{i=0}^t c_i x_i$, with $A(x) = B(x) \cdot C(x)$, and recall the induced matrix factorization $\llbracket A \rrbracket_\ell = \llbracket B \rrbracket_\ell \cdot \llbracket C \rrbracket_{\ell+s}$. To prove that

$$\pi_\ell \circ \llbracket A \rrbracket_\ell = \pi_\ell \circ \llbracket B \rrbracket_\ell \circ \llbracket C \rrbracket_{\ell+s}$$

is surjective from $I_{\epsilon, m}$ to $\mathbb{T}^\ell$, we can just prove that $\pi_\ell \circ \llbracket B \rrbracket_\ell$ is surjective from $I_{\delta, \ell+s}$ for some δ , and that $\llbracket C \rrbracket_{\ell+s} \cdot I_{\epsilon, m}$ contains $I_{\delta, \ell+s}$. We anticipate that $C(x)$ will be choosed monic and with all roots < 1 in absolute value.

3.1. Estimate for $B(x)$. We can take $\delta = 1/b_0$. Let's show that the image of the cube $I_{\delta, \ell+s}$ under $\llbracket B \rrbracket_\ell$ assumes each value modulo $\mathbb{Z}^\ell$, starting with an arbitrary $v = (v_1, \dots, v_\ell) \in \mathbb{R}^\ell$ and building inductively a vector $w = (w_1, \dots, w_{\ell+s})$ in $I_{\delta, \ell+s}$ such that $\llbracket B \rrbracket_\ell \cdot w - v \in \mathbb{Z}^\ell$. Suppose that $w \in I_{\delta, \ell+s}$ is such that the components with index $> i$ of $\llbracket B \rrbracket_\ell \cdot w - v$ are in $\mathbb{Z}$, and observe that while w_i varies in the interval $[-\delta/2, \delta/2]$ the i -th component of $\llbracket B \rrbracket_\ell \cdot w$ varies in an interval large $b_0\delta = 1$, while all components with index $> i$ of $\llbracket B \rrbracket_\ell \cdot w$ stay fixed. Consequently we can move w_i in the interval $[-\delta/2, \delta/2]$ to ensure that all components with index $\geq i$ of $\llbracket B \rrbracket_\ell \cdot w - v$ are in $\mathbb{Z}$, and since repeating this procedure allows us to construct a w with the required properties, our claim follows.

3.2. Estimate for $C(x)$. If $C(x) = \sum_{i=0}^t c_i x^i = \prod_{i=1}^t (x - \gamma_i)$, we will see that it is possible to take

$$\epsilon = \delta \cdot \prod_{i=1}^t \frac{1}{1 - |\gamma_i|}.$$

To prove the assertion, we proceed in the following way. Rather than working with $\llbracket C \rrbracket_{\ell+s}$ we will work with the nonsingular square matrix $\{C\}_m$, and if we prove that the image under $\{C\}_m$ of $I_{\epsilon, m}$ contains $I_{\delta, m}$ our claim will follow, since the image under $\llbracket C \rrbracket_m$ is a just projection on the last $\ell + s$ coordinates of the image under $\{C\}_m$.

Now, rather than proving that the image under $\{C\}_m$ of $I_{\epsilon, m}$ contains $I_{\delta, m}$, it will be easier to prove that the image under $\{C\}_m^{-1}$ of $I_{\delta, m}$ is contained in $I_{\epsilon, m}$. In particular, $\{C\}_m$ factors as

$$\{C\}_m = \prod_{i=1}^t \{x - \gamma_i\}_m,$$

and the inverse of a matrix of the form $\{x - \gamma\}_m$, for $\gamma \in \{\gamma_i\}_{1 \leq i \leq t}$, is easily computed as

$$\begin{pmatrix} 1 & & & & & & \\ -\gamma & 1 & & & & & \\ & -\gamma & 1 & & & & \\ & & -\gamma & \ddots & & & \\ & & & \ddots & 1 & & \\ & & & & -\gamma & 1 & \end{pmatrix}^{-1} = \begin{pmatrix} 1 & & & & & & \\ \gamma & 1 & & & & & \\ \gamma^2 & \gamma & 1 & & & & \\ \gamma^3 & \gamma^2 & \gamma & \ddots & & & \\ \vdots & \vdots & \vdots & \ddots & 1 & & \\ \gamma^{m-1} & \gamma^{m-2} & \dots & \gamma^2 & \gamma & 1 & \end{pmatrix}.$$

This shows that if a (possibly complex) vector $v = (v_1, \dots, v_m)$ has all components with absolute value $\leq \rho$ for some real number $\rho > 0$, the vector w obtained applying the matrix $\{x - \gamma\}_m^{-1}$ will have components of the form

$$w_{r+1} = \sum_{i=0}^r \gamma^i v_{r+1-i}$$

for some $0 \leq r < m$, and their absolute value can be estimated as

$$\left| \sum_{i=0}^r \gamma^i v_{r+1-i} \right| \leq \sum_{i=0}^r |\gamma^i| \cdot |v_{r+1-i}| \leq \sum_{i=0}^r |\gamma^i| \cdot \rho \leq \frac{1}{1-|\gamma|} \rho.$$

Since ρ is arbitrary we have applying iteratively $\{x - \gamma_i\}_m^{-1}$ for $i = 1, \dots, t$ that the set of complex vectors with all components $< \delta$ in absolute value is mapped by $\{C\}_m^{-1}$ to complex vectors with components of absolute value at most ϵ , and consequently $I_{\delta,m}$ is mapped into $I_{\epsilon,m}$, being $\{C\}_m^{-1}$ a matrix with real entries.

3.3. Conclusion. Let $A(x) = B(x)C(x)$ be a real factorization of $A(x)$, with

$$B(x) = \sum_{i=0}^s b_i x^i = a_d \cdot \prod_{i=1}^s (x - \beta_i), \quad C(x) = \sum_{i=0}^t c_i x^i = \prod_{i=1}^t (x - \gamma_i),$$

and with $C(x)$ monic with all roots < 1 in absolute value. The above estimate shows that Q is ϵ -dense for each ϵ at least

$$\frac{1}{|a_d|} \cdot \prod_{i=0}^s \frac{1}{|\beta_i|} \cdot \prod_{i=1}^t \frac{1}{1 - |\gamma_i|},$$

and consequently for ϵ at least

$$\frac{1}{|a_d|} \cdot \prod_{i=0}^d \min \left\{ \frac{1}{|\alpha_i|}, \frac{1}{1 - |\alpha_i|} \right\} = \left(|a_d| \cdot \prod_{i=1}^d \max \left\{ |\alpha_i|, 1 - |\alpha_i| \right\} \right)^{-1}$$

since we can take as γ_i precisely the α_i with absolute value $\leq 1/2$, and as β_i the remaining roots of $A(x)$ (note that $C(x)$ will have real coefficients). Since this last expression is clearly not greater than $1/M(A(x/2))$, the proof is complete. $\square$

4. INTEGRAL LINEAR RECURRENCES OF FIXED LENGTH


Let $A(x) = \sum_{i=0}^d a_i x^i$ be a primitive integral polynomial of degree d with $a_0 \neq 0$, and for $m > d$ let Λ_m be the $\mathbb{Z}$ -module of the integral vectors in $\mathbb{Z}^m$ that form a recurrence sequence determined by $A(x)$. It is also the module of the integral vectors that are in the kernel of the matrix $\llbracket A \rrbracket_{m-d}$, and as such has rank d . In this section we provide some results about the structure of this $\mathbb{Z}$ -module, and in particular we show that for each prime p we can choose a basis with special properties with respect to the p -adic valuation of the vectors.

Only a small amount of these results is required to prove Theorem 2, but we will state and prove them in full strength nevertheless. We will consider the $\mathbb{Z}_p$ -module $\Lambda_m^{(p)} = \Lambda_m \otimes_{\mathbb{Z}} \mathbb{Z}_p$ over the ring of p -adic integers $\mathbb{Z}_p$ for simplicity, which is also clearly equal to the vectors in $\mathbb{Z}_p^m$ killed by $\llbracket A \rrbracket_{m-d}$, having $\llbracket A \rrbracket_{m-d}$ integral coefficients. All considerations can then be lifted to $\mathbb{Z}$ with arbitrary approximation with respect to the p -adic absolute value.

Let $v_p(\cdot)$ be the p -adic valuation, and let's recall that the Newton polygon of $A(x)$ is the boundary of the polygon in $\mathbb{R}^2$ obtained as the upper convex envelop of the points with coordinates

$$(0, v_p(a_0)), (1, v_p(a_1)), \dots, (d, v_p(a_d)).$$

Suppose that the polygon has r edges with different slopes which connect the consecutive pairs of points $(w_k, v_p(a_{w_k}))$ for $0 = w_0 < w_1 < \dots < w_r = d$, let $\ell_k = w_k - w_{k-1}$ be the length of the horizontal projection of the k -th edge, and let $\sigma_k = (v_p(a_{w_k}) - v_p(a_{w_{k-1}}))/\ell_k$ be its slope. We have that $\sigma_i < \sigma_j$ whenever $1 \leq i < j \leq d$.



Newton polygon of $A(x) = 9x^4 - 3x^3 - 9x^2 - 2x + 3$ for $p = 3$.

Let's consider a splitting field K of $A(x)$ over the field of p -adic numbers $\mathbb{Q}_p$, and let's extend to K the p -adic valuation and absolute value. Call as usual $\mathcal{O}_K$ the integral closure of $\mathbb{Z}_p$ in K , which is the local ring formed by the element with non-negative valuation, with maximal ideal $\mathfrak{m}$ generated by the uniformizer π . As it is well known, the polynomial $A(x)$ has precisely ℓ_k roots with p -adic valuation equal to $-\sigma_k$, for $1 \leq k \leq r$. We will show that the slopes of the Newton polygon, and consequently the valuations of the roots of $A(x)$, are reflected in the $\mathbb{Z}_p$ -module $\Lambda_m^{(p)}$ as stated in the following:

Theorem 3. *For each prime p there exist a unique basis of $\Lambda_m^{(p)}$ such that if we form the $d \times m$ matrix $M = (M_{i,j})$ with the basis vectors as row vectors, then*

- (1) *the left $d \times d$ submatrix $(M_{i,j})_{\substack{1 \leq i \leq d \\ 1 \leq j \leq d}}$ (resp. the submatrix $(M_{i,j})_{\substack{1 \leq i \leq d \\ m-d+1 \leq j \leq m}}$) is a block upper (resp. lower) triangular matrix, with blocks $B_1, \dots, B_r$ (resp. $C_1, \dots, C_r$) on the diagonal; the number of blocks r is equal to the number of edges of the Newton polygon of $A(x)$, and B_k and C_k are square matrices of size equal to the length ℓ_k of the k -th side of the polygon for each $1 \leq k \leq r$;*
- (2) *if $1 \leq s \leq r$ is the smallest integer such that $\sigma_s \geq 0$ (or we may require alternatively that s is the smallest such that $\sigma_s > 0$) then $B_1, \dots, B_{s-1}, C_s, \dots, C_r$ are the identity matrix, and for $1 \leq k \leq s-1$ the matrix C_k has determinant with valuation $-\sigma_k \ell_k (m-d)$, and for $s \leq k \leq r$ the matrix B_k has determinant with valuation $\sigma_k \ell_k (m-d)$;*
- (3) *walking on the right from the element $M_{j,j} = 1$ contained in the diagonal of the k -th identity block B_k for $1 \leq k \leq s-1$ (resp. walking left from an $M_{j+m-d,j} = 1$ contained in the diagonal of the*

identity block C_k , for $s \leq k \leq r$) by t steps we find elements with p -adic valuation at least $-\tau\sigma_k$ (resp. at least $\tau\sigma_k$).

In other words, the matrix M can be taken with the following form:

$$\begin{pmatrix} \boxed{1} & & \dots & & \dots & \boxed{C_0} & & & & \\ & \boxed{1} & & \dots & & \dots & \boxed{C_1} & & & 0 \\ & & \ddots & & & \dots & \ddots & & & \\ & & & \boxed{1} & \dots & \dots & \dots & \boxed{C_{s-1}} & & \\ & & & & \boxed{B_s} & \dots & \dots & & \boxed{1} & \\ & & & & & \ddots & & & & \ddots \\ & 0 & & & & & \boxed{B_{r-1}} & \dots & & \dots & \boxed{1} \\ & & & & & & & \boxed{B_r} & \dots & & \boxed{1} \end{pmatrix}$$

and the behaviour of the p -adic valuation on the rows is controlled by the slope of the segments of the Newton polygon of $A(x)$.

For instance, if $A(x) = 9x^4 - 3x^3 - 9x^2 - 2x + 3$ and we take $p = 3$, then for recurrence length $m = 10$ the matrix M is

$$\begin{pmatrix} \boxed{1} & \frac{480}{887} & \frac{4203}{16853} & \frac{3861}{33706} & \frac{2511}{33706} & \frac{243}{16853} & \boxed{\frac{729}{33706}} & 0 & 0 & 0 \\ 0 & \frac{-7722}{887} & \frac{-5049}{16853} & \frac{-3339}{33706} & \frac{-76419}{33706} & \frac{33378}{16853} & \frac{-51543}{33706} & 1 & 0 & 0 \\ 0 & \frac{729}{887} & \frac{-44658}{16853} & \frac{42822}{16853} & \frac{-27306}{16853} & \frac{19179}{16853} & \frac{3489}{16853} & 0 & 1 & 0 \\ 0 & 0 & 0 & \boxed{\frac{729}{1078}} & \frac{243}{1078} & \frac{405}{539} & \frac{675}{1078} & \frac{423}{539} & 48/49 & \boxed{1} \end{pmatrix}.$$

All denominators are prime with 3, the 3-adic valuations of the elements are

$$\begin{pmatrix} 0 & 1 & 2 & 3 & 4 & 5 & 6 & \infty & \infty & \infty \\ \infty & 3 & 3 & 2 & 2 & 1 & 3 & 0 & \infty & \infty \\ \infty & 6 & 3 & 3 & 2 & 2 & 1 & \infty & 0 & \infty \\ \infty & \infty & \infty & 6 & 5 & 4 & 3 & 2 & 1 & 0 \end{pmatrix},$$

and as it is possible observe the valuation increases going right in the first row, slowly increases going left in the second and third rows, and increases going left on the last row.

Proof. Let $N = (N_{i,j})$ be the rational $d \times m$ matrix obtained putting $N_{i,j} = \delta_{ij}$ for $1 \leq i, j \leq d$, and inductively defining the remaining elements in each row to form a linear recurrence determined by $A(x)$. We will now prove that the square matrix $N_\xi = (N_{i,\xi_j})_{1 \leq i, j \leq d}$ is nonsingular for $\xi = (1, 2, \dots, w, m - d + w + 1, \dots, m)$, for all $w = w_0, \dots, w_r$ which are the ordinate of a vertex of the Newton polygon of $A(x)$.

If $A(x)$ has distinct roots $\alpha_1, \dots, \alpha_d$ the matrix N is given by $V^{-1} \cdot L$, where

$$V = (\alpha_i^{j-1})_{1 \leq i, j \leq d}, \quad L = (\alpha_i^{j-1})_{\substack{1 \leq i \leq d \\ 1 \leq j \leq m}}.$$

To obtain a formula for the determinant of N_ξ valid for general α_i , let's work over $\mathbb{C}$ and suppose for a moment that $\frac{1}{a_d}A(x) = \prod_{i=1}^d (x - \alpha_i)$ where the α_i are algebraically independent over $\mathbb{C}$. The determinant of $N_\xi = (N_{i,\xi_j})_{1 \leq i, j \leq d}$

is equal to the determinant of $L_\xi = (L_{i,\xi_j})_{1 \leq i,j \leq d}$ divided by $\det V$, and this turns out to be the Schur function s_λ (see [10]) associated to the partition λ defined as $\lambda_{d-1+i} = \xi_i - i + 1$ for $1 \leq i \leq d$, evaluated in $\alpha_1, \dots, \alpha_d$. Note applying the definition of N by linear recurrence that the entries of N are polynomial functions in the elementary symmetric functions of the α_i , and hence they are polynomial in the α_i , and the determinant of each submatrix is polynomial in the α_i as well. While the α_i vary outside of the closed algebraic set defined by $\prod_{i>j} (\alpha_i - \alpha_j) = 0$ the determinant of N_ξ is equal to the polynomial $s_\lambda(\alpha_1, \dots, \alpha_d)$, and consequently the equality must hold for each value of the α_i .

If ξ is defined as above, λ has precisely $d-w$ parts all equal to $m-d$, and its conjugate partition λ' is formed by $m-d$ parts equal to $d-w$. Recall now Jacobi-Trudi's identity

$$s_\lambda = \det(e_{\lambda'_i - i + j})_{1 \leq i,j \leq k},$$

where the e_i are the elementary symmetric functions, and which holds for each k at least as big as the number of parts of λ' . We are reduced to prove that a matrix of the form $(e_{d-w-i+j})_{1 \leq i,j \leq m-d}$ is non-singular. After evaluation in the roots we have $e_i = (-1)^i a_{d-i}/a_d$, and flipping the sign of rows and columns of even order we can consider the determinant of

$$U = (a_{w+i-j})_{1 \leq i,j \leq m-d}$$

up to possibly a sign, and discarding a factor a_d^{-m+d} . Note that all entries on the diagonal are equal to a_w .

If $w = w_0 = 0$ (resp. if $w = w_r = d$) then the matrix is lower (resp. upper) triangular with $a_w \neq 0$ on the diagonal, and consequently non-singular. Suppose $w = w_k$ for some $1 < k < r$, let $\rho \in K$ be an element with p -adic valuation equal to σ_k , and let R be the diagonal matrix with $1, \rho, \rho^2, \dots, \rho^{m-d-1}$ on the diagonal. The matrix

$$\frac{1}{a_w} R^{-1} \cdot U \cdot R = \left(\frac{a_{w+i-j} \rho^{j-i}}{a_w} \right)_{1 \leq i,j \leq m-d}$$

has all entries in $\mathcal{O}_K$, and is upper unitriangular when reduced modulo $\mathfrak{m}$ because it has all 1 on the diagonal, and starting from a_w the p -adic valuation increases at a rate bigger than σ_k .

Consequently N_ξ is non-singular, and we can consider the matrix $Q = N_\xi^{-1} \cdot N$. This is already sufficient to show that the matrix M must be unique, because suppose that we have another matrix M' with the same properties, and suppose that the i -th row differs. If $(v_j)_{1 \leq j \leq m}$ is the difference of those rows we must have $v_j = 0$ unless $w_k < j \leq m-d+w_k$ for some k , so taking $w = w_k$ we have that (v_j) cannot be a linear combination of the rows of Q . But Q has rank d , and the existence of (v_j) would imply that the module of linear recurrences $\Lambda_m \otimes_{\mathbb{Z}} \mathbb{Q}$ has rank $> d$, which is absurd.

Suppose now that $w = w_k$ with $1 \leq k < s$, so that the slope σ_k is ≤ 0 by definition of s . We will show that all entries in the rows with indices $w_{k-1} + 1, \dots, w_k - 1, w_k$ are integral, and that the valuation of $Q_{i,j}$ for $w_{k-1} < i \leq w_k$ and $j \geq i$ is at least $-\sigma_k(j-i)$.

In fact, let ρ have valuation equal to σ_k and consider the polynomial

$$B(x) = \sum_{i=0}^d \frac{\rho^{w-i} a_i}{a_w} x^i = \sum_{i=0}^d b_i x^i \in \mathcal{O}_K[x].$$

Note that the Newton polygon of $B(x)$ is obtained by the Newton polygon of $A(x)$ by subtracting the linear affine function $f(x) = \sigma_k(x - w) + v_p(a_w)$, and this has the effect of moving the k -th side of the Newton polygon to lay on the horizontal axis. In particular, the coefficients of $B(x)$ are such that $b_{w_{k-1}}$ and b_{w_k} are $\not\equiv 0 \pmod{\mathfrak{m}}$, but $b_i \equiv 0 \pmod{\mathfrak{m}}$ for $i < w_{k-1}$ or $i > w_k$.

On the other hand, for each $w_{k-1} < i \leq w_k$ the vector $(\rho^{j-i} Q_{i,j})_{1 \leq j \leq m}$ is a linear recurrence determined by $B(x)$, and we claim that all entries are in $\mathcal{O}_K$. In fact, suppose this is not the case, and multiply it by the smallest power of the uniformizer π required to make all entries in $\mathcal{O}_K$. Some entry will be in $\mathcal{O}_K \setminus \mathfrak{m}$, but the first $\ell_k = w_k$ entries will be in $\mathfrak{m}$. When reduced modulo $\mathfrak{m}$, the subvector $(\rho^{j-i} Q_{i,j})_{w_{k-1} < j \leq m-d+w_k}$ is a recurrence determined by the polynomial

$$C(x) = \sum_{i=0}^{\ell_k} \overline{b_{w_{k-1}+i}} x^i = \sum_{i=0}^{\ell_k} c_i x^i \in (\mathcal{O}_K/\mathfrak{m})[x],$$

and this recurrence of order ℓ_k in $\mathcal{O}_K/\mathfrak{m}$ is supposed to have non-zero entries, while the first ℓ_k entries are zero. But this is absurd, and the claim is proved.

The matrix $(\rho^{j-i} Q_{i,j})_{\substack{w_{k-1} < i \leq w_k \\ m-d+w_{k-1} < j \leq m-d+w_k}}$ is invertible, because modulo $\mathfrak{m}$ it is obtained as the $(m-d)$ -th power of the companion matrix

$$\begin{pmatrix} 0 & & & -c_0/c_{\ell_k} \\ 1 & 0 & & -c_1/c_{\ell_k} \\ & 1 & 0 & -c_2/c_{\ell_k} \\ & & \ddots & \vdots \\ & & & 1 & -c_{\ell_k-1}/c_{\ell_k} \end{pmatrix}$$

of $C(x)$, which is invertible. Consequently $(Q_{i,j})_{\substack{w_{k-1} < i \leq w_k \\ m-d+w_{k-1} < j \leq m-d+w_k}}$, which

is obtained by it conjugating by the diagonal matrix with diagonal $1, \rho, \dots, \rho^{\ell_k-1}$ and multiplying by a factor $\rho^{-(m-d)}$, has the same valuation as $\rho^{-\ell_k(m-d)}$, i.e. equal to $-\sigma_k \ell_k (m-d)$.

In this way we have built the rows from $w_{k-1} + 1$ to w_k of M , and proved that $C_k = (Q_{i,j})_{\substack{w_{k-1} < i \leq w_k \\ m-d+w_{k-1} < j \leq m-d+w_k}}$ has determinant with valuation $-\sigma_k \ell_k (m-d)$, while $B_k = (Q_{i,j})_{w_{k-1} < i, j \leq w_k}$ is the identity. For $s \leq k \leq r$ we can clearly proceed in a symmetrical way, taking C_k equal to the identity and proceeding on the left up to B_k which will have determinant equal to $\sigma_k \ell_k (m-d)$.

The matrix we have built selecting at each step the rows from $w_{k-1} + 1$ to w_k of Q clearly satisfies all requirements for the matrix M . Furthermore all rows are linearly independent, so the module they generate over $\mathbb{Z}_p$ has rank d .

To prove that the rows generate all of $\Lambda_m^{(p)}$ observe that they generate a $\mathbb{Z}_p$ -module contained in $\Lambda_m^{(p)}$, and suppose the generated module to be strictly contained. By left multiplication by a matrix B with determinant $\in \mathbb{Q}_p \setminus \mathbb{Z}_p$ we can obtain in any case a basis of $\Lambda_m^{(p)}$, but the matrix $M_\xi = (M_{i,\xi_j})_{1 \leq i,j \leq d}$ for $\xi = (1, \dots, w_s, m - d + w_s + 1, \dots, m)$ has determinant 1, and $B \cdot M_\xi$ (and consequently $B \cdot M$) would not have coefficients in $\mathbb{Z}_p$, which is absurd. $\square$

Let's now turn to the $\mathbb{Z}$ -module Λ_m again. If E is a $\mathbb{Z}$ -module and $F \subseteq E$ a submodule with index n , we have that the index of $F \otimes_{\mathbb{Z}} \mathbb{Z}_p$ in $E \otimes_{\mathbb{Z}} \mathbb{Z}_p$ is precisely the biggest power of p dividing n .

Let $\Theta_m \subseteq \mathbb{Z}^d \times \mathbb{Q}^{m-d}$ be the $\mathbb{Z}$ -module of the linear recurrences determined by $A(x)$ such that the first d coordinates are in $\mathbb{Z}$, and similarly let $\Theta_m^{(p)} = \Theta_m \otimes_{\mathbb{Z}} \mathbb{Z}_p$. The matrix N of the proof of the Theorem is clearly a $\mathbb{Z}$ -basis of Θ_m .

For each prime p the matrix M is equal to $M_\xi \cdot N$ where $M_\xi = (M_{i,\xi_j})_{1 \leq i,j \leq d}$ and $\xi = (1, 2, \dots, d)$, and as it is possible to verify immediately M_ξ has determinant with the same p -adic valuation as a_d^{m-d} , because for $s \leq k \leq r$ the matrix B_k has determinant with valuation equal to $\sigma_k \ell_k(m - d) = (v_p(a_{w_k}) - v_p(a_{w_{k-1}}))(m - d)$.

Consequently $(\Theta_m^{(p)} : \Lambda_m^{(p)})$ is equal to the biggest power of p dividing a_d^{m-d} , for each p . We have proved:

Corollary 1. *The module Λ_m has index equal to $|a_d|^{m-d}$ in Θ_m .*

5. PROOF OF THEOREM 2

Before providing the proof of Theorem 2, let's recall a few facts of linear algebra. Let $\langle \cdot, \cdot \rangle$ be the standard inner product on $\mathbb{R}^n$, and $u_1, \dots, u_k \in \mathbb{R}^n$, for $k \leq n$. The Gram matrix of the u_i is the $k \times k$ matrix defined as

$$G(u_1, \dots, u_k) = (\langle u_i, u_j \rangle)_{1 \leq i,j \leq k},$$

and geometrically its determinant is the square of the volume of the parallelepiped formed by the vectors u_i . If the u_i can be completed with $u_{k+1}, \dots, u_n \in \mathbb{R}^n$ to a basis of $\mathbb{R}^n$, and $v_1, v_2, \dots, v_n \in \mathbb{R}^n$ are such that they form a pair of biorthonormal bases, then we have

$$G(u_1, u_2, \dots, u_n)G(v_1, v_2, \dots, v_n) = I,$$

(see [8, §66.2, pag. 66-6]). Since $G(u_1, \dots, u_k)$ is a minor of $G(u_1, u_2, \dots, u_n)$ and its complementary cofactor is $G(v_{k+1}, \dots, v_n)$, if the $u_1, \dots, u_n$ form a parallelepiped of volume 1 (and consequently $G(u_1, \dots, u_n)$ has determinant 1) we have that

$$\det G(u_1, \dots, u_k) = \det G(v_{k+1}, \dots, v_n),$$

as it follows from the properties of compound matrices (see [6, Chap. 1, §4, pag. 21, equation (33) in particular]) related to what sometimes is also called "Jacobi's Theorem" [7, §14.16].

It is now possible to give the

Proof of Theorem 2. As in the proof of Theorem 1, put $\ell = m - d$, and let Q be defined as

$$Q = Q(A) = \left\{ v \in \mathbb{R}^m : \llbracket A \rrbracket_\ell \cdot v \in \mathbb{Z}^\ell \right\}.$$

Let $\omega_1, \dots, \omega_d \in \mathbb{Z}^m$ be any basis of the d -dimensional the lattice Λ_m of vectors in $\mathbb{Z}^m$ which are killed by $\llbracket A \rrbracket_\ell$ (i.e. form a linear recurrence determined by $A(x)$), and let $e_1, \dots, e_m$ be the standard basis of $\mathbb{R}^m$.

Suppose by contradiction $\bar{\epsilon} < \epsilon < 1/M(A(x))$ are numbers, independent of m , such that Q is $\bar{\epsilon}$ -dense. Since $\epsilon > \bar{\epsilon}$ we have $\pi_m(Q + [0, \epsilon]^m) = \mathbb{T}^m$, and since Q is the union of integral translates of the parallelepiped formed by $\omega_1, \dots, \omega_d$ the map $\pi_m : \mathbb{R}^m \rightarrow \mathbb{T}^m$ must be surjective on the parallelepiped Π formed by combinations with coefficients in $[0, 1]$ of the vectors $\epsilon e_1, \dots, \epsilon e_m, \omega_1, \dots, \omega_d$:

$$\Pi = \left\{ \sum_{i=1}^m d_i \cdot \epsilon e_i + \sum_{i=1}^d c_i \cdot \omega_i, \quad c_i, d_i \in [0, 1] \right\}.$$

The map π_m locally preserves the volume and the image of Π is all $\mathbb{T}^m$, so the volume of Π must be ≥ 1 . On the other side, the volume of Π can be computed as the sum of the volumes of the parallelepiped formed by all choices of m vectors within $\epsilon e_1, \dots, \epsilon e_m, \omega_1, \dots, \omega_d$, and note that the volume of the parallelepiped formed by, say, $\epsilon e_{s_1}, \dots, \epsilon e_{s_q}, \omega_{r_1}, \dots, \omega_{r_p}$ with $p + q = m$ is not greater than

$$\epsilon^q \sqrt{G(\omega_{r_1}, \omega_{r_2}, \dots, \omega_{r_p})} \leq \epsilon^{m-d} \sqrt{G(\omega_{r_1}, \omega_{r_2}, \dots, \omega_{r_p})},$$

being $q \geq m - d$ and $\epsilon \leq 1$. On the other side the total number of such parallelepipeds is $\binom{m+d}{m}$.

The theorem now follows if we prove that we can select a special basis $\omega_1, \dots, \omega_d$ of the lattice Λ to ensure that $G(\omega_{r_1}, \dots, \omega_{r_p}) < K(M)M^{2m}$ for each subset of the ω_i and for each $M > M(A)$, for some constant $K(M)$ not dependent of m . In fact, volume of Π must clearly be

$$\text{Vol}(\Pi) \leq \sqrt{K(M)} \binom{m+d}{m} \epsilon^{m-d} M^m$$

and in particular it $\rightarrow 0$ as $m \rightarrow \infty$ if we take $M(A) < M < 1/\epsilon$, which we can do since we supposed $\epsilon < 1/M(A)$.

To construct the required basis, let $\zeta_1, \dots, \zeta_d \in \mathbb{Q}^m$ be such that the j -th coordinate of ζ_i is δ_{ij} for $1 \leq i, j \leq d$, and defining the remaining coordinates by the linear recurrence determined by $A(x)$. The ζ_i are clearly a basis of the $\mathbb{Z}$ -module Θ_m of vectors in $\mathbb{Z}^d \times \mathbb{Q}^\ell$ which are killed by $\llbracket A \rrbracket_\ell$, and we have by Corollary 1 that $(\Lambda_m : \Theta_m) = |a_d|^\ell$. Consequently, a basis (ω_i) of Λ_m can be obtained by the (ζ_i) applying a matrix $W = (W_{i,j})_{1 \leq i, j \leq d}$ with determinant $|a_d|^\ell$. Changing it by left multiplication with an element of $SL(d, \mathbb{Z})$ the matrix W can be supposed to be in Hermite Normal Form [8, §23.2, pag. 23-6, 23-7], and in particular upper triangular and such that $|W_{i,j}| \leq |W_{j,j}|$ whenever $i < j$.

We will now show that we can bound the Gramian of a subset of the ω_i in terms of the Gramians of all subsets of the ζ_i and the determinant of $(W_{i,j})$ squared (which we know to be equal to $|a_d|^{2\ell}$). In fact, let $\omega_{r_1}, \dots, \omega_{r_p}$, for $1 \leq p \leq d$ and $1 \leq r_1 < \dots < r_p \leq d$, be a subset of the ω_i . The quantity $\sqrt{G(\omega_{r_1}, \dots, \omega_{r_p})}$, being the volume of the parallelepiped formed by the ω_{r_i} , is also equal to

$$\sup_{\phi \in \Phi_p} \phi(\omega_{r_1}, \dots, \omega_{r_p}),$$

the sup being taking withing all the elements of the exterior power $\Lambda_p^*(\mathbb{R}^m)$ of the form

$$\Phi_p = \{n_1 \wedge n_2 \wedge \dots \wedge n_p, \text{ for orthonormal } n_1, \dots, n_p \in \mathbb{R}^m\} \subset \Lambda_p^*(\mathbb{R}^m),$$

where $\mathbb{R}^m$ is identified with the dual through $\langle \cdot, \cdot \rangle$. This can now be estimated as

$$\begin{aligned} & \sup_{\phi \in \Phi_p} \phi\left(\sum_{j=1}^d W_{r_1,j} \zeta_j, \dots, \sum_{j=1}^d W_{r_p,j} \zeta_j\right) \\ &= \sup_{\phi \in \Phi_p} \sum_{1 \leq s_1, \dots, s_p \leq d} \phi(W_{r_1,s_1} \zeta_{s_1}, \dots, W_{r_p,s_p} \zeta_{s_p}) \\ &\leq \sum_{1 \leq s_1, \dots, s_p \leq d} \sup_{\phi \in \Phi_p} W_{r_1,s_1} \cdots W_{r_p,s_p} \cdot \phi(\zeta_{s_1}, \dots, \zeta_{s_p}) \\ &\leq \sum_{1 \leq s_1, \dots, s_p \leq d} |W_{s_1,s_1} \cdots W_{s_p,s_p}| \cdot \sup_{\phi \in \Phi_p} \phi(\zeta_{s_1}, \dots, \zeta_{s_p}) \\ &\leq \det(W_{i,j}) \cdot \sum_{1 \leq s_1, \dots, s_p \leq d} \sup_{\phi \in \Phi_p} \phi(\zeta_{s_1}, \dots, \zeta_{s_p}), \end{aligned}$$

since we can discard the summands where $s_i = s_j$ for some $i \neq j$, and $(W_{i,j})$ is upper triangular and with integral entries,

$$\leq \det(W_{i,j}) \cdot d^p \cdot \max_{1 \leq s_1 < \dots < s_p \leq d} \sqrt{G(\zeta_{s_1}, \dots, \zeta_{s_p})}.$$

Note in particular that, for each $M > M(A(x))$, this allows us to convert an estimate of the form

$$G(\zeta_{r_1}, \dots, \zeta_{r_p}) < K(M) [|a_d^{-1}| \cdot M]^{2m}, \quad \text{for all } 1 \leq r_1 < \dots < r_p \leq d \quad (1)$$

into one of the form

$$G(\omega_{r_1}, \dots, \omega_{r_p}) < K'(M) M^{2m}, \quad \text{for all } 1 \leq r_1 < \dots < r_p \leq d$$

for some other constant $K'(M) = d^p K(M)$ (thus uniformly in m).

To prove the (1), let $B(x) = a_d^{-1} A(x)$ (so that $M(B) = |a_d^{-1}| \cdot M(A)$), and let $(B_i)_{1 \leq i \leq \ell}$ be the row vectors of the rational matrix $\llbracket B \rrbracket_\ell$. Note that it can be completed to a matrix with determinant 1 inserting the row vectors $e_1, \dots, e_d$ of the standard basis $(e_i)_{1 \leq i \leq m}$ of $\mathbb{R}^m$, and the ζ_i are dual to the e_i in the basis $e_1, \dots, e_d, B_1, \dots, B_\ell$ with respect to the standard scalar product $\langle \cdot, \cdot \rangle$.

Consequently, let $\zeta_{r_1}, \dots, \zeta_{r_p}$ for $1 \leq r_1 < \dots < r_p \leq d$ be a subset of the ζ_i , and let $1 \leq s_1 < \dots < s_q \leq d$ be the complementary set of the r_i in $1, 2, \dots, d$. As observed before the proof, the Gram determinant of $\zeta_{r_1}, \dots, \zeta_{r_p}$ is the same as the Gram determinant of the vectors $e_{s_1}, \dots, e_{s_q}, B_1, \dots, B_\ell$. Note that this Gram matrix is a matrix with a very special form, and after a complicated part we have that if j, k are $\geq q$ the $(j+1, k+1)$ entry is equal to the (j, k) entry being $\langle B_{j-q}, B_{k-q} \rangle = \langle B_{j+1-q}, B_{k+1-q} \rangle$, and in particular it is a perturbed Toeplitz matrix.

The asymptotic behaviour of matrices that are perturbed Toeplitz matrices have been deeply studied by Bump-Diaconis [2], Tracy-Widom [12] and Lyons [9]. In particular, the following theorem of Lyons [9, Theorem 3.1] suits perfectly our needs.

Let λ the Lebesgue measure on the unit circle, and let $E_n(t) = e^{2\pi i n t}$. Let Poly_n be the linear span of $E_0, E_1, \dots, E_n$. Let $H^2(\mathbb{T})$ be the Hardy space of the functions $f \in L^2(\lambda)$ such that all negative Fourier coefficients vanish. Let P_{H^2} be the orthogonal projection $L^2(\lambda) \rightarrow H^2(\mathbb{T})$. For $f \geq 0$ with $\log f \in L^1(\lambda)$, let

$$\Phi_f(z) = \exp \frac{1}{2} \int_{\mathbb{T}} \frac{E_1(t) + z}{E_1(t) - z} \log f(t) d\lambda(t),$$

for $|z| < 1$. Let $\phi_f(t)$ be the *outer* function

$$\phi_f(t) := \lim_{r \uparrow 1} \Phi_f(rE_1(t)),$$

which exists λ -almost everywhere [11], and satisfies $|\phi_f|^2 = f$, λ -almost everywhere. If μ is a measure on $\mathbb{T}$, let its Fourier coefficients be $\hat{\mu}(n) = \int_{\mathbb{T}} e^{-2\pi i n t} d\mu(t)$, and let D_n be the determinant of the Toeplitz matrix $(\hat{\mu}(k-j))_{0 \leq j, k \leq n}$. Then we have the following result:

Theorem 4 (R. Lyons). *Let μ be a finite positive measure on $\mathbb{T}$ with infinite support. Let $f := [d\mu/d\lambda]$ be the Radon-Nikodým derivative of the absolutely continuous part of μ .*

Given any functions $f_0, \dots, f_r, g_0, \dots, g_r \in L^2(\mu)$, let $F_j := P_{H^2}(f_j \overline{\phi_f})$ and $G_j := P_{H^2}(g_j \overline{\phi_f})$. Define

$$P_j := \begin{cases} f_j & \text{if } 0 \leq j \leq r, \\ E_j & \text{if } -1 \geq j \geq -n \end{cases}$$

and

$$Q_j := \begin{cases} g_j & \text{if } 0 \leq j \leq r, \\ E_j & \text{if } -1 \geq j \geq -n. \end{cases}$$

We have

$$\lim_{n \rightarrow \infty} D_{n-1}(\mu)^{-1} \det \left[\int P_j \overline{Q_k} d\mu \right]_{-n \leq j, k \leq r} = \det \left[\int F_j \overline{G_k} d\lambda \right]_{0 \leq j, k \leq r}.$$

Note that we can take $n = \ell, r = q-1, \mu = f\lambda$ for $f = B(e^{2\pi i t})B(e^{-2\pi i t})$, and $f_j = g_j = e^{-2\pi i s_{j+1}t} B(e^{-2\pi i t})^{-1}$, which clearly $\in L^2(\mu)$, for $0 \leq j \leq q-1$.

In this way after a trivial computation we get

$$\left[\int P_j \overline{Q_k} d\mu \right]_{-n \leq j, k \leq r} = G(e_{s_1}, \dots, e_{s_q}, B_1, \dots, B_\ell),$$

because for instance for $1 \leq j \leq q$ and $1 \leq k \leq \ell$ we have

$$\begin{aligned} \int P_{j-1} \overline{Q_{-k}} d\mu &= \int_0^1 e^{-2\pi i s_j t} B(e^{-2\pi i t})^{-1} \cdot e^{2\pi i k t} \cdot B(e^{2\pi i t}) B(e^{-2\pi i t}) dt \\ &= \int_0^1 e^{2\pi i (-s_j + k)t} \cdot B(e^{2\pi i t}) dt \\ &= b_{s_j - k} = \langle e_{s_j}, B_k \rangle. \end{aligned}$$

On the other hand $(\hat{\mu}(k - j))_{0 \leq j, k \leq n}$ after is the Toeplitz matrix associated to the Laurent polynomial $B(x)B(x^{-1})$ as can be readily verified, and consequently $D_{n-1}(\mu) = G(B_1, \dots, B_\ell)$.

The theorem tells us that for $\ell \rightarrow \infty$ the asymptotic value of the ratio

$$\det G(e_{s_1}, \dots, e_{s_q}, B_1, \dots, B_\ell) / \det G(B_1, \dots, B_\ell)$$

is finite for each $s_1, \dots, s_q \in \{1, \dots, d\}$, so we are reduced to show that

$$\det G(B_1, \dots, B_\ell) \leq K(M)M^{2m}$$

for each $M > M(B)$ for some constant $K(M)$.

But a nice result on the determinant of banded Hermitian Toeplitz matrices is available, thanks to the Trench's Formula [1, Theorem 2.10, pag. 41]: let $C(x) = \sum_{j=-r}^s c_j x^j$ be a Laurent polynomial, and let $g_n(z)$ be the row

$$g_n(z) = (1, z, z^2, \dots, z^{r-1}, z^{n+r}, z^{n+r+1}, \dots, z^{n+r+s-1}).$$

Let $\xi_1, \dots, \xi_k$ be the distinct roots of $C(x)$, and let $\mu_1, \dots, \mu_k$ be their multiplicities. We define G_n as the determinant of the $(r+s) \times (r+s)$ matrix Γ_n whose first μ_1 rows are $g_n(\xi_1), g'_n(\xi_1), \dots, g_n^{(\mu_1-1)}(\xi_1)$, whose next μ_2 rows are $g_n(\xi_2), g'_n(\xi_2), \dots, g_n^{(\mu_2-1)}(\xi_2)$, and so on.

Theorem 5 (Trench). *We have $G_0 \neq 0$, and*

$$D_{n-1}(C) = (-1)^{ns} c_s^n \frac{G_n}{G_0}, \quad \text{for every } n \geq 1.$$

Applying the Theorem with $C(x) = B(x)B(x^{-1})$ and $n = \ell$, we just have to show that $c_s^n G_n$ is smaller than $K(M)M^n$ for each $M > M(C) = M(B)^2$, for some $K(M)$. But if we expand the determinant of the $(r+s) \times (r+s)$ matrix Γ_n we have that it can be written as sum of monomials in the ξ_i with polynomials in n as coefficients, where in each monomial ξ_i appears with exponent smaller than $(n+r+s)\mu_i$. Consequently estimating each monomial the determinant is $\leq P(n)M(\frac{1}{c_s}C)^{n+r+s}$ for some polynomial $P(n)$ in n , and in this way we obtain an estimate of the form $c_s^n G_n \leq K(M)M^n$ for each $M > M(C)$, as required. $\square$

REFERENCES

- [1] A. Böttcher and S.M. Grudsky. *Spectral Properties of Banded Toeplitz Matrices*. Society for Industrial and Applied Mathematics Philadelphia, PA, USA, 2005.
- [2] D. Bump and P. Diaconis. Toeplitz minors. *Journal of Combinatorial Theory, Series A*, 97(2):252–271, 2002.
- [3] J.W.S. Cassels. An introduction to diophantine approximation, Cambridge Tracts 45, 1957.
- [4] J.W.S. Cassels. An introduction to the geometry of numbers. 2nd printing, corrected. 1971.
- [5] G. Everest and T. Ward. *Heights of polynomials and entropy in algebraic dynamics*. Springer Verlag, 1999.
- [6] F.R. Gantmacher. *The theory of matrices, Vol. 1*. Chelsea Pub. Co., New York, 1959.
- [7] I.S. Gradshteyn, I.M. Ryzhik, A. Jeffrey, and D. Zwillinger. *Table of integrals, series, and products*. Academic Press, 2000.
- [8] L. Hogben, R.A. Brualdi, A. Greenbaum, and R. Mathias. *Handbook of linear algebra*. CRC Press, 2007.
- [9] R. Lyons. Szego limit theorems. *Geometric And Functional Analysis*, 13(3):574–590, 2003.
- [10] I.G. Macdonald. *Symmetric Functions and Hall Polynomials (2nd edition)*. Oxford University Press, New York, 1995.
- [11] W. Rudin. *Real and complex analysis*. Tata McGraw-Hill, 2006.
- [12] C.A. Tracy and H. Widom. On the limit of some Toeplitz-like determinants. *SIAM Journal on Matrix Analysis and Applications*, 23(4):1194–1198, 2002.

DIPARTIMENTO DI MATEMATICA "L. TONELLI" - LARGO BRUNO PONTECORVO, 5
- 56127 PISA

E-mail address: dubbini@mail.dm.unipi.it

SCUOLA NORMALE SUPERIORE DI PISA - PIAZZA DEI CAVALIERI, 7 - 56126 PISA

E-mail address: maurizio.monge@sns.it